

Final approval of the article: MM, JM, HB, JP, AL, SR, LH, CK

Statistical analysis: JM

Obtained funding: MM

Overall responsibility: MM

## REFERENCES

1. Kearon C, Kahn S, Agnelli G, Goldhaber S, Raskob GE, Comerota AJ. Antithrombotic therapy for venous thromboembolic disease: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines (8th edition). *Chest* 2008;301:454-545S.
2. Lagerstedt CI, Olsson CG, Fagher BO, Oqvist BW, Albrechtsson U. Need for long-term anticoagulant treatment in symptomatic calf-vein thrombosis. *Lancet* 1985;2:515-8.
3. Hull R, Delmore T, Genton E, Hirsh J, Gent M, Sackett D, et al. Warfarin sodium versus low-dose heparin in the long-term treatment of venous thrombosis. *N Engl J Med* 1979;301:855-8.
4. Levine MN, Hirsh J, Gent M, Turpie AG, Weitz J, Ginsberg J, et al. Optimal duration of oral anticoagulant therapy: a randomized trial comparing four weeks with three months of warfarin in patients with proximal deep vein thrombosis. *Thromb Haemost* 1995;74:606-11.
5. Kearon C, Ginsberg JS, Kovacs MJ, Anderson DR, Wells P, Julian JA, et al. Extended Low-Intensity Anticoagulation for Thromboembolism Investigators. Comparison of low-intensity warfarin therapy with conventional-intensity warfarin therapy for long-term prevention of recurrent venous thromboembolism. *N Engl J Med* 2003;349:631-9.
6. Ridker PM, Goldhaber SZ, Danielson E, Rosenberg Y, Eby CS, Deitcher SR, et al. PREVENT Investigators. Long-term, low-intensity warfarin therapy for the prevention of recurrent venous thromboembolism. *N Engl J Med* 2003;348:1425-34.
7. Piazza G, Seddighzadeh A, Goldhaber SZ. Deep-vein thrombosis in the elderly. *Clin Appl Thromb Hemost* 2008;14:393-8.
8. Spencer FA, Gore JM, Lessard D, Emery C, Pacifico L, Reed G, et al. Venous thromboembolism in the elderly. A community-based perspective. *Thromb Haemost* 2008;100:780-8.
9. Arcelus JJ, Caprini JA, Monreal M, Suárez C, González-Fajardo J. The management and outcome of acute venous thromboembolism: a prospective registry including 4011 patients. *J Vasc Surg* 2003;38:916-22.
10. Laporte S, Mismetti P, Découssus H, Uresandi F, Otero R, Lobo JL, et al. RIETE investigators. Clinical predictors for fatal pulmonary embolism in 15,520 patients with venous thromboembolism; findings from the Registro Informatizado de la Enfermedad TromboEmbolica venosa (RIETE) Registry. *Circulation* 2008;117:1711-6.
11. Trujillo-Santos J, Nieto JA, Tiberio G, Piccoli A, di Micco P, Prandoni P, et al. RIETE Registry. Predicting recurrences or major bleeding in cancer patients with venous thromboembolism. Findings from the RIETE registry. *Thromb Haemost* 2008;100:435-9.
12. López-Jiménez L, Montero M, González-Fajardo JA, Arcelus JJ, Suárez C, Lobo JL, et al, for the RIETE investigators. Venous thromboembolism in very elderly patients: findings from a prospective registry (RIETE). *Haematologica* 2006;96:1046-51.
13. Vasco B, Villalba JC, Lopez-Jimenez L, Falgá C, Montes J, Trujillo-Santos J, et al, for the RIETE Investigators. Venous thromboembolism in nonagenarians. Findings from the RIETE Registry. *Thromb Haemost* 2009;101:1112-8.
14. Linkins LA, Choi PT, Douketis JD. Clinical impact of bleeding in patients taking oral anticoagulant therapy for venous thromboembolism: a meta-analysis. *Ann Intern Med* 2003;139:893-900.
15. Kearon C, Ginsberg JS, Julian JA, Douketis J, Solymoss S, Ockelford P, et al. Fixed-dose heparin (FIDO) Investigators. Comparison of fixed-dose weight-adjusted unfractionated heparin and low-molecular-weight heparin for acute treatment of venous thromboembolism. *JAMA* 2006;296:935-42.
16. Schulman S, Kearon C, Kakkar AK, Mismetti P, Schellong S, Eriksson H, et al. RE-COVER Study group. Dabigatran versus warfarin in the treatment of acute venous thromboembolism. *N Engl J Med* 2009;361:2342-52.
17. Carrier M, Le Gal G, Wells PS, Rodger MA. Systematic review: case-fatality rates of recurrent venous thromboembolism and major bleeding events among patients treated for venous thromboembolism. *Ann Intern Med* 2010;152:578-89.

Submitted Apr 26, 2011; accepted May 27, 2011.

## APPENDIX

### Members of the RIETE group

Coordinator of the RIETE Registry: Dr Manuel Monreal (Spain)

RIETE Steering Committee Members: Dr Hervé Decousus (France), Dr Paolo Prandoni (Italy), and Dr Benjamin Brenner (Israel)

RIETE National Coordinators: Dr Raquel Barba (Spain), Dr Pierpaolo Di Micco (Italy), and Dr Karine Rivron-Guillot (France)

RIETE Registry Coordinating Center: S & H Medical Science Service

SPAIN: Arcelus JJ, Barba R, Blanco A, Barrón M, Casado I, Casas JM, Cisneros E, del Campo R, del Toro J, Durán M, Falgá C, Fernández-Capitán C, Gabriel F, Gallego P, García-Bragado F, Gómez-Zorilla S, Guijarro R, Guil M, Gutiérrez J, Hernández L, Hernández-Huerta D, Jiménez D, Jiménez M, Jordán S, Lecumberri R, Lobo JL, López L, Lorenzo A, Luque JM, Madridano O, Maestre A, Marchena PJ, Martín-Villasclaras JJ, Monreal M, Nauffal MD, Nieto JA, Núñez MJ, Oribe M, Otero, R, Pedrajas JM, Rabuñal R, Riera-Mestre A, Román P, Rosa V, Rubio S, Ruiz FJ, Ruiz-Giménez N, Sahuquillo JC, Samperiz A, Sánchez Muñoz-Torrero JF, Sánchez R, Soler S, Soler C, Tiberio G, Todolí JA, Tolosa C, Trujillo J, Uresandi F, Valdés V, Valle R, Vela J, Vidal G, Villalta J.

FRANCE: Boccalon H, Delluc A, Farge-Bancel D, Mahe I, Rivron-Guillot K. ISRAEL: Brenner B.

ITALY: Barillari A, Barillari G, Ciammaichella M, Dalla Valle F, Di Micco P, Duce R, Piovella C, Poggio R, Prandoni P, Pasca S, Quintavalla R, Schenone A, Tiraferri E, Visonà A.

REPUBLIC of MACEDONIA: Bosevski M.

SWITZERLAND: Bounameaux H.

## INVITED COMMENTARY

**Anthony J. Comerota, MD, Toledo, Ohio; and Ann Arbor, Mich**

Sanchez Munoz-Torrero and the RIETE investigators report that older patients with acute deep vein thrombosis (DVT) have a higher mortality from bleeding than pulmonary embolism (PE) in the first 3 months. Although point estimates are low, their obser-

vations may be surprising to most readers. Their findings reaffirm prior observations that major bleeding events from anticoagulation, particularly vitamin K antagonists (VKAs), occur early in the course of therapy.

This robust database has provided enormous insight into the outcome of patients with DVT and PE. Most recurrences follow the initial event, that is, patients presenting with PE will have PE as a recurrent event, and those presenting with DVT recur with DVT. That was also the case in these patients.

The authors also showed that the diagnosis of DVT carries an early mortality ranging from 6% to 11% in patients  $\geq 60$  years of age. The majority of the fatalities that were observed occurred beyond the first week of therapy; therefore, one would assume that the majority of the patients were being treated with a VKA. Two factors on multivariate analysis were found to correlate with fatal PE: chronic lung disease and malignancy. However, eight factors were associated with fatal bleeding: age (upper decades), low body weight, recent major bleeding, renal dysfunction, bilateral DVT, cancer, immobility  $\geq 4$  days, and prior venous thromboembolism. Aside from renal dysfunction, low body weight, and recent major bleeding, it is difficult to integrate the bleeding risk factors to modify clinical care. The authors tabulated outcome events by decade of life in Table II. The most impressive discrepancy in mortality occurred in the patients  $\geq 80$  years of age, where major/fatal bleeding occurred in 3.6% vs thrombotic events in 1.7%. Eleven percent of this age group died within this 3-month window.

It is reasonable to assume that all major bleeding will be clinically apparent and therefore accurately recognized. However,

that is not the case with fatal PE. To fulfill the definition of fatal PE, the patients must have experienced an objectively confirmed PE followed by death within 10 days. Therefore, sudden unexplained deaths, which more commonly would occur out of the hospital, would not be recorded as a fatal PE by the authors' definition. As they recognized, the rate of fatal PE may be underestimated. This is not to detract from the value of this report nor the importance of the RIETE registry.

The authors observe that in older patients, the risk of fatal bleeding exceeds the risk of fatal PE in patients presenting with acute DVT. Clinicians might assume that less aggressive anticoagulation therapy with VKAs might be indicated. Unfortunately, that cannot be substantiated by observations from a prior randomized trial.<sup>1</sup> It will be interesting to observe how guideline writing committees respond to the important observations reported by the RIETE investigators.

## REFERENCE

1. Kearon C, Ginsberg JS, Kovacs MJ, Anderson DR, Wells P, Julian JA, et al. Comparison of low-intensity warfarin therapy with conventional-intensity warfarin therapy for long-term prevention of recurrent venous thromboembolism. *N Engl J Med* 2003;349:631-9.